

Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at <u>www.bccancer.bc.ca</u> and according to acceptable standards of care

PROTOCOL CODE: BRLACTWACG

(Page 1 of 2)

DOCTOR'S ORDERS Htcm Kg BSA	m²		
REMINDER: Please ensure drug allergies and previous bleomycin are documented on the	Allergy & Alert Form		
DATE: To be given: Cycle #:			
Date of Previous Cycle:			
Delay treatment week(s)			
CBC & Diff, platelets day of treatment			
For PAClitaxel and CARBOplatin (Days 1, 8, and 15): May proceed with doses as written if within 48 h ANC greater than or equal to 1.5 x 10 ⁹ /L, Plate equal to 90 x 10 ⁹ /L	elets <u>greater than or</u>		
For DOXOrubicin and cyclophosphamide:			
May proceed with doses as written if within 72 h ANC greater than or equal to 1.0 x 10 ⁹ /L, Platelets greater than or equal to 100 x 10 ⁹ /L			
Dose modification for: Imatology Other Toxicity			
Proceed with treatment based on blood work from			
PREMEDICATIONS: Patient to take own supply. RN/Pharmacist to confirm	·		
45 Minutes Prior to PACLitaxel: dexamethasone 10 mg IV in NS 50 mL over 15 minutes 30 Minutes Prior to PACLitaxel: diphenhydrAMINE 25 mg IV in NS 50 mL over 15 minutes and famotidine 20 mg IV in NS 100 mL over 15 minutes (Y-site compatible) ondansetron 8 mg PO 30 minutes prior to CARBOplatin			
No pre-medication required (see protocol for guidelines)			
OR dexamethasone 3 mg or 12 mg (select one) PO 30 to 60 minutes prior to AC treatment and select ONE of the following:			
ondansetron 8 mg PO 30 to 60 minutes prior to AC treatment			
aprepitant 125 mg PO 30 to 60 minutes prior to AC treatment			
 ondansetron 8 mg PO 30 to 60 minutes prior to AC treatment netupitant-palonosetron 300 mg-0.5 mg PO 30 to 60 minutes prior to AC treatment 			
Other:			
Have Hypersensitivity Reaction Tray and Protocol Available for Cycles	1 to 4		
CHEMOTHERAPY: (Note – continued over 2 pages)			
☐ CYCLE # (Cycle 1-4)			
PACLitaxel 80 mg/m² OR mg/m² (select one) x BSA = mg Dose Modification: % = mg/m² x BSA = mg IV in 100 to 500 mL (non-DEHP bag) NS over 1 hour on Days 1, 8 and 15 (use non-DEHP tubing with 0.2 micron in-			
line filter)			
CARBOplatin AUC □ 6 or □ 5 or □ 4 (select one) x (GFR + 25) = mg □ Dose Modification: % = mg IV in 100 to 250 mL NS over 30 minutes on Day 1			
*** SEE PAGE 2 FOR CHEMOTHERAPY CYCLES 5 TO 8 ***			
DOCTOR'S SIGNATURE:	SIGNATURE:		
	UC:		
BC Cancer Provincial Preprinted Order BRLACTWACG			



Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at <u>www.bccancer.bc.ca</u> and according to acceptable standards of care

PROTOCOL CODE: BRLACTWACG

(Page 2 of 2)

DOCTOR'S ORDERS	Page 2 of 2		
DATE:	To be given:	Cycle #:	
CHEMOTHERAPY continued			
CYCLE # (Cycle 5-	-8)		
DOXOrubicin 60 mg/m ² x BSA = Dose Modification: IV push	mg % = mg/m² x BSA :	= mg	
cyclophosphamide 600 mg/m ² x BS/ Dose Modification: IV in 100 to 250 mL NS over 20 m	% = mg/m² x BSA :	= mg	
RETURN APPOINTMENT ORDERS			
 Return in two or three weeks for D x 3 for cycles 1-4; book chemo room e 13) 	octor and Cycle every two weeks for AC cycles	_ (Book chemo room weekly 5-8, cycle 5 to start week	
☐ Post Cycle 5: Book filgrastim (G-C	SF) SC teaching and first dos	e on Day:	
Last Cycle. Return in	week(s) after last treat	ment.	
Cycles 1 to 4: CBC & Diff, Platelets, Creatinine prior to each cycle. CBC & Diff, Platelets prior to treatment on days 8 and 15.			
<u>Cycles 5 to 8:</u> CBC & Diff, Platelets prior to each cy	/cle.		
If clinically indicated: 🗌 ALT 🔲 Bilirubin 🗌 MUGA 🗌 Echocardiogram			
☐ Other tests:			
☐ Consults:			
See general orders sheet for ad	ditional requests.		
DOCTOR'S SIGNATURE:			SIGNATURE:
			UC: