

Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at <a href="www.bccancer.bc.ca">www.bccancer.bc.ca</a> and according to acceptable standards of care

## PROTOCOL CODE: UGIFFOXPAN

(Page 1 of 2)

BC Cancer "Compassionate Access Program" approval must be obtained prior to treatment

DOCTOR'S ORDERS	Ht	cm	Wt	kg	BSA_	m²
REMINDER: Please ensure drug allerg	ies and previous ble	eomyci	n are do	ocumented	on the A	Allergy & Alert Form
DATE:	To be given:			Cycle	e(s) #:	
Date of Previous Cycle:						
Delay treatment week(s) CBC & Diff, Platelets day of treatment May proceed with doses as written if within 72 hours ANC greater than or equal to 1.2 x 109/L, Platelets greater than or equal to 75 x 109/L Dose modification for:						
PREMEDICATIONS: Patient to take own supply. RN/Pharmacist to confirm  ondansetron 8 mg PO prior to treatment dexamethasone ☐ 8 mg or ☐ 12 mg (select one) PO prior to treatment NO ice chips ☐ Other:						
magnesium sulfate 2 g in 50 mL NS over 30 min for hypomagnesemia magnesium sulfate 5 g in 100 mL NS over 3 hours for hypomagnesemia						
**Have Hyper	sensitivity Reaction	Tray a	nd Prot	ocol Availal	le**	
CHEMOTHERAPY: (Note – continued over 2 pages)  Repeat in two weeks Repeat in two and in four weeks  PANitumumab 6 mg/kg x kg = mg  Dose Modification:mg/kg x kg = mg  IV in 100 mL NS over 1 hour. If tolerated, administer over 30 minutes in subsequent cycles. Use 0.2 micron in-line filter. Flush lines with 25 mL NS pre and post PANitumumab infusion.						
Prior to starting oxaliplatin, flush lines  oxaliplatin 85 mg/m² x BSA =mg  Dose Modification:mg  IV in 250 to 500 mL D5W over 2 hours	mg	mg				
☐ leucovorin 400 mg/m² x BSA = mg IV in 250 mL D5W over 2 hours*  * oxaliplatin and leucovorin may be infused over same two hour period by using a Y-site connector placed immediately before the injection site.						
OR						
☐ leucovorin 20 mg/m² x BSA = mg IV push  *** SEE PAGE 2 FOR FLUOROURACIL CHEMOTHERAPY ***						
DOCTOR'S SIGNATURE:					SIG UC:	NATURE:



Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at <a href="www.bccancer.bc.ca">www.bccancer.bc.ca</a> and according to acceptable standards of care

## PROTOCOL CODE: UGIFFOXPAN

(Page 2 of 2)

DATE:							
CHEMOTHERAPY: (Continued)							
fluorouracil 400 mg/m² x BSA = mg							
	Dose Modification:mg/m² x BSA =mg						
IV push							
fluorouracil 2400 mg/m <sup>2</sup> x BSA = mg**							
☐ Dose Modification:mg/m² x BSA =mg**							
IV over 46 hours in D5W to a total volume of 230 mL by continuous infusion at 5 mL/h via Baxter LV5 INFUSOR							
	** For 3000 to 5500 mg dose, <b>select INFUSOR per dose range below (doses outside</b> dose banding range are						
prepared as ordered):							
Doco Panding Pange	Pharmaniat II	nitial and Date					
Dose Banding Range Less than 3000 mg	Dose Band INFUSOR (mg)  Pharmacy to mix specific dose	Pharmacist	illiai and Date				
3000 to 3400 mg	3200 mg						
3401 to 3800 mg	3600 mg						
3801 to 4200 mg	4000 mg						
4201 to 4600 mg	4400 mg						
4601 to 5000 mg	4800 mg						
5001 to 5500 mg	5250 mg						
Greater than 5500 mg	Pharmacy to mix specific dose						
	·						
	RETURN APPOINTMEN	T ORDERS					
Return in two weeks for Doo	tor and Cycle						
, <del></del>							
Return in <u>four</u> weeks for Do							
Return in <u>six</u> weeks for Doc							
□ Loot Cyclo Detumaia							
Last Cycle. Return in							
CBC & Diff, Platelets, Creatini							
prior to each cycle							
☐ INR weekly ☐ INR prior to							
☐ ECG ☐ CEA ☐ Other tests:							
☐ Book for PICC assessmen							
☐ Book for IVAD insertion pe							
☐ Weekly Nursing Assessme							
☐ Consults:							
☐ See general orders sheet f							
DOCTOR'S SIGNATURE:			SIGNATURE:				
			uc.				
			UC:				