

Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at www.bccancer.bc.ca and according to acceptable standards of care

PROTOCOL CODE: UMYBLDF

Page 1 of 2

Patient RevAid ID: A BC Cancer "Compassionate Access Program" request form must be completed and approved prior to treatment. **DOCTOR'S ORDERS** m^2 Ht cm Wt kg BSA REMINDER: Please ensure drug allergies and previous bleomycin are documented on the Allergy & Alert Form DATE: To be given: Cycle #: Date of Previous Cycle: Risk Category: Female of Childbearing Potential (FCBP) Rx valid for 7 days Risk Category: Male or Female of non-Childbearing Potential (NCBP) ____ week(s) ☐ Delay treatment ☐ CBC and Diff, Platelets, Creatinine, ALT, Bilirubin on day of treatment ☐ CBC on day of treatment May proceed with lenalidomide dose day 1 as written, if within 96 hours ANC greater than or equal to 1.0 x 10⁹/L, platelets greater than or equal to 50 x 10⁹/L and eGFR as per protocol May proceed with bortezomib dose day 1 as written, if within 96 hours ANC greater than or equal to 0.5 x 10⁹/L, platelets greater than or equal to 50 x 10⁹/L. bilirubin less than or equal to 1.5 x upper limit of normal • If CBC prior to day 1 show ANC less than 1.5 x 109/L or platelets less than 75 x 109/L then: May proceed with bortezomib Day 8 and 15 as written, if within 96 hours ANC greater than or equal to 0.5 x 10⁹/L. platelets greater than or equal to 50 x 109/L Dose modification for: Hematology Renal Function Other Toxicity Proceed with treatment based on blood work from Pharmacy Use for Lenalidomide **LENALIDOMIDE** dispensing: One cycle = 28 days ☐ lenalidomide* _____mg PO daily, in the evening, on days 1 to 21 and off for 7 days ☐ lenalidomide* _____ mg PO ______(*a Part Fill # 1 (*available as 25 mg, 20 mg, 15 mg, 10 mg, 5 mg, 2.5 mg capsules) RevAid confirmation number: *Note: Use one capsule strength for the total dose; there are cost implications as costing Lenalidomide lot number: is per capsule and not weight based ☐ FCBP dispense 21 capsules (1 cycle) Pharmacist counsel (initial): ☐ For Male and Female NCBP: capsules or cycles. Maximum 63 capsules (3 cycles). Pharmacy to Part Fill # 2 dispense one cycle at a time, maximum 3 cycles if needed STEROID*: CHOOSE ONE RevAid confirmation number: One cycle = 28 days ☐ dexamethasone ☐ 40 mg or ☐ 20 mg PO once weekly, in the morning, Lenalidomide lot number: x _____ doses **OR** __number of 28 day cycles_____(select one) Pharmacist counsel (initial): mg PO once weekly in the morning, x ☐ doses dexamethasone OR ☐ number of 28 day cycles_____ (select one) predniSONE _____mg PO once weekly in the morning, x _____doses OR Part Fill # 3 number of 28 day cycles____ (select one) RevAid confirmation number: ☐ No Steroid *Refer to Protocol for steroid dosing options Lenalidomide lot number: Physician to ensure DVT prophylaxis in place: ☐ ASA, ☐ Warfarin, ☐ low molecular Pharmacist counsel (initial): weight heparin, \(\square\) direct oral anticoagulant or \(\subseteq \) none (select one) **Special Instructions DOCTOR'S SIGNATURE:** SIGNATURE: Physician RevAid ID: UC:

Created: 1 Feb 2021 Revised: 1 April 2021 (formatting revised)



Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at www.bccancer.bc.ca and according to acceptable standards of care

PROTOCOL CODE: UMYBLDF

Page 2 of 2

DATE:	
TREATMENT : If patient is VZV seropositive and/or at physician's clinical judgement, physician to ensure prophylaxis with valACYclovir 500 mg daily while on bortezomib and for 4 weeks after discontinuation	
CYCLE # (Cycles 1 to 8)	
bortezomib _ 1.3 mg /m² or _ 1 mg/m² or _ 0.7 mg/m² (select one) x BSA = mg SC injection on days 1, 8, and 15	
RETURN APPOINTMENT ORDERS	
For Cycles 1 to 8, book chemo on Days 1, 8 and 15	
Return in four weeks for Doctor and Cycle	
Last cycle. Return inweek(s)	
Laboratory: Blood work done prior to next cycle must be done less than or equal to 4 days prior to the start date	
TSH every three months (i.e. prior to cycles 1, 4, 7, 10,13 etc)	
Cycles 1 to 4: CBC and Diff, Platelets, Creatinine, Calcium every two weeks	
Day 1: CBC and Diff, Platelets, Creatinine, Calcium, bilirubin, ALT Serum Protein Electrophoresis <u>and/or</u> Serum Free Light Chain Levels (SELECT APPROPRIATE)	
CBC and Diff, Platelets on Day 8 and 15 for current cycle if ANC on Day 1 is less than $1.5 x$ $10^9/L$ or Platelets are less than $100 x$ $10^9/L$	
Cycles 5 to 8: Day 1: CBC and Diff, Platelets, Creatinine, Calcium, bilirubin, ALT ☐ Serum Protein Electrophoresis and/or ☐ Serum Free Light Chain Levels (SELECT APPROPRIATE)	
CBC and Diff, Platelets on Day 8 and 15 for current cycle if ANC on Day 1 is less than 1.5 x $10^9/L$ or Platelets are less than 75 x $10^9/L$	
<u>Cycle 9 and subsequent cycles:</u> Day 1: CBC and Diff, Platelets, Creatinine, Calcium, bilirubin, ALT ☐ Serum Protein Electrophoresis <u>and/or</u> ☐ Serum Free Light Chain Levels (SELECT APPROPRIATE)	
☐ Pregnancy blood test for FCBP 7-14 days and 24 h prior to cycle 1 and every week for 4 weeks during cycle 1	
☐ Pregnancy blood test for FCBP, every 4 weeks, less than or equal to 7 days prior to the next cycle	
☐ Other tests ☐ Consults:	
☐ See general orders sheet for additional requests	
DOCTOR'S SIGNATURE:	SIGNATURE: UC: