

Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at www.bccancer.bc.ca and according to acceptable standards of care

PROTOCOL CODE: SAAVGEMD

Page 1 of 1

To continue beyond 6 cycles, a BC Cancer "Compassionate Access Program" request must be approved.

DOCTOR'S ORDERS Htcm Wt	kg BSAm²
REMINDER: Please ensure drug allergies and previous bleomycin are document	
DATE: To be given:	Cycle #:
Date of Previous Cycle:	
□ Delay Treatment week(s) □ CBC & Diff, Platelets day of treatment May proceed with doses as written Day 1 if within 24 hours ANC greater than or equal to 0.9 x 10 ⁹ /L and Platelets greater than or equal to 100 x 10 ⁹ /L May proceed with doses as written Day 8 if within 24 hours ANC greater than or equal to 1.4 x 10 ⁹ /L and Platelets greater than or equal to 100 x 10 ⁹ /L Dose modification for: □ Hematology □ Other Toxicity	
Proceed with treatment based on bloodwork from	
PREMEDICATIONS: Patient to take own supply. RN/Pharmacist to confirm dexamethasone 8 mg PO BID for 3 days, starting one day prior to each DOCEtaxel treatment. (Note: A minimum of 3 doses of dexamethasone pre-treatment is required.) ondansetron 8 mg PO prior to treatment prochlorperazine 10 mg PO or ☐ metoclopramide 10 mg (select one) PO prior to treatment Optional: Frozen gloves starting 15 minutes before DOCEtaxel infusion until 15 minutes after end of DOCEtaxel infusion; gloves should be changed after 45 minutes of wearing. Other:	
Have Hypersensitivity Reaction Tray and Protocol Available	
CHEMOTHERAPY: gemcitabine	
, , , , , , , , , , , , , , , , , , ,	
RETURN APPOINTMEN Return in three weeks for Doctor and Cycle Book Cher Last Cycle. Return inweeks.	
CBC & Diff, Platelets prior to each treatment (Day 1 and 8) Prior to Cycle 4: Bilirubin, LDH ALT, GGT, Alk Phos If clinically indicated:	
DOCTOR'S SIGNATURE:	SIGNATURE:
	UC: RN: