

1-, 3- and 5-year Observed Survival and Age-Standardized Relative Survival (Table 1) and associated 95% Confidence Intervals (Table 2), by Stage at Diagnosis for Breast, Colorectal (Colon, Rectum), Prostate and Lung, 2017*

Table 1: Survival Estimates

| ne 1. Sur vivar Es | | | | | Age-standardized Relative Survival | | | |
|--------------------|--------------|--------|--------------------------|--------|------------------------------------|--------------|--------|--|
| | | | Observed Survival Rate % | | | Rate Ratio % | | |
| Cancer Type | Cancer Stage | 1 Year | 3 Year | 5 Year | 1 Year | 3 Year | 5 Year | |
| Breast (Female) | Stage I | 99.0 | 96.0 | 92.8 | 100.5 | 100.5 | 100.9 | |
| | Stage II | 97.9 | 91.8 | 84.5 | 99.6 | 96.5 | 91.6 | |
| | Stage III | 94.6 | 79.5 | 67.3 | 95.5 | 82.0 | 70.6 | |
| | Stage IV | 67.6 | 38.9 | 22.2 | 68.4 | 40.4 | 24.0 | |
| | Stage UNK | 75.2 | 60.1 | 51.3 | 82.5 | 73.5 | 68.2 | |
| Colorectal | Stage I | 96.3 | 90.8 | 84.0 | 98.6 | 98.5 | 97.7 | |
| | Stage II | 91.9 | 82.1 | 72.5 | 95.9 | 91.8 | 88.0 | |
| | Stage III | 89.8 | 73.5 | 62.7 | 92.7 | 80.2 | 72.9 | |
| | Stage IV | 49.3 | 19.9 | 11.9 | 53.0 | 22.5 | 14.2 | |
| | Stage UNK | 70.0 | 56.5 | 48.9 | 81.3 | 73.2 | 69.1 | |
| Colon | Stage I | 95.8 | 90.2 | 83.3 | 98.4 | 98.1 | 97.3 | |
| | Stage II | 91.7 | 82.4 | 72.7 | 96.1 | 92.9 | 89.7 | |
| | Stage III | 87.9 | 71.2 | 61.4 | 91.8 | 79.6 | 73.5 | |
| | Stage IV | 47.5 | 18.6 | 11.1 | 51.9 | 21.6 | 13.6 | |
| | Stage UNK | 64.7 | 52.6 | 44.8 | 79.1 | 72.3 | 68.5 | |
| Rectum | Stage I | 97.0 | 90.9 | 84.3 | 98.9 | 97.8 | 96.3 | |
| | Stage II | 92.5 | 80.1 | 70.8 | 95.2 | 86.6 | 80.9 | |
| | Stage III | 93.5 | 78.3 | 65.4 | 94.6 | 81.9 | 72.0 | |
| | Stage IV | 56.5 | 24.2 | 14.4 | 57.5 | 25.2 | 15.4 | |
| | Stage UNK | 84.2 | 68.3 | 59.9 | 88.7 | 78.2 | 72.9 | |
| Lung | Stage I | 87.3 | 66.6 | 53.2 | 91.4 | 76.4 | 66.3 | |
| | Stage II | 70.5 | 44.8 | 34.7 | 77.7 | 54.2 | 45.2 | |
| | Stage III | 50.6 | 21.8 | 15.1 | 57.0 | 28.7 | 22.6 | |
| | Stage IV | 23.1 | 7.4 | 4.4 | 27.8 | 9.5 | 5.7 | |
| | Stage UNK | 35.5 | 20.6 | 16.3 | 48.1 | 36.9 | 31.3 | |
| Prostate | Stage I | 97.3 | 92.7 | 89.6 | 99.3 | 98.7 | 100.4 | |
| | Stage II | 97.8 | 92.8 | 87.8 | 100.3 | 100.6 | 101.1 | |
| | Stage III | 99.1 | 95.6 | 90.6 | 101.0 | 100.1 | 101.2 | |
| | Stage IV | 80.5 | 52.2 | 37.8 | 87.0 | 63.3 | 49.6 | |
| | Stage UNK | 86.5 | 72.8 | 64.1 | 95.0 | 90.4 | 87.8 | |

NOTE: * The end year of the most recent period that varies for different cancer types. -- Not applicable. If the standard error of a survival rate is between 0.05 and 0.1, the 95% confidence interval of the rate is shown in the table. If it is > 0.1 or is not estimable due to a small number of cases or events then the survival rate is suppressed and denotes as §. Survival rates include age at diagnosis 15 - 99.

Please refer to the <u>Introduction and Technical Notes</u> for further information.

Data Source: BC Cancer Registry Date Retrieved: 17DEC2019 Analysis by: Cancer Surveillance and Outcomes, Data & Analytics



Table 2: 95% Confidence Intervals (CI)

| 2. 93 /0 Con | | | | | 95% CI of Age-standardized Relative | | | |
|-----------------|--------------|-------------|------------------------------------|-------------|-------------------------------------|-----------------------|--------------|--|
| | | 95% CI of O | 95% CI of Observed Survival Rate % | | | Survival Rate Ratio % | | |
| Cancer Type | Cancer Stage | 1 Year | 3 Year | 5 Year | 1 Year | 3 Year | 5 Year | |
| Breast (Female) | Stage I | 98.7 - 99.2 | 95.5 - 96.5 | 92.1 - 93.5 | 100.1 - 100.8 | 99.7 - 101.2 | 99.7 - 102.0 | |
| | Stage II | 97.5 - 98.3 | 91.0 - 92.5 | 83.4 - 85.6 | 99.0 - 100.1 | 95.4 - 97.6 | 89.9 - 93.3 | |
| | Stage III | 93.3 - 95.6 | 77.5 - 81.4 | 64.9 - 69.5 | 93.9 - 96.8 | 79.3 - 84.5 | 67.2 - 73.9 | |
| | Stage IV | 64.0 - 70.9 | 35.3 - 42.5 | 19.0 - 25.5 | 64.8 - 71.7 | 36.5 - 44.4 | 20.1 - 28.2 | |
| | Stage UNK | 70.1 - 79.5 | 54.9 - 64.9 | 46.4 - 56.0 | 77.9 - 86.3 | 68.0 - 78.4 | 62.1 - 73.7 | |
| Colorectal | Stage I | 95.5 - 97.0 | 89.6 - 91.9 | 82.4 - 85.5 | 97.8 - 99.3 | 97.2 - 99.7 | 95.7 - 99.6 | |
| | Stage II | 90.8 - 92.8 | 80.6 - 83.4 | 70.8 - 74.1 | 95.0 - 96.7 | 90.3 - 93.2 | 85.9 - 90.0 | |
| | Stage III | 88.7 - 90.7 | 72.0 - 74.9 | 61.0 - 64.4 | 91.7 - 93.6 | 78.6 - 81.8 | 70.7 - 74.9 | |
| | Stage IV | 47.2 - 51.4 | 18.3 - 21.5 | 10.6 - 13.3 | 50.9 - 55.0 | 20.7 - 24.3 | 12.6 - 15.9 | |
| | Stage UNK | 67.5 - 72.4 | 53.8 - 59.1 | 46.2 - 51.5 | 79.3 - 83.1 | 70.6 - 75.6 | 66.1 - 71.9 | |
| Colon | Stage I | 94.9 - 96.6 | 88.8 - 91.5 | 81.3 - 85.0 | 97.4 - 99.1 | 96.5 - 99.5 | 94.9 - 99.4 | |
| | Stage II | 90.5 - 92.8 | 80.8 - 84.0 | 70.8 - 74.5 | 95.0 - 96.9 | 91.2 - 94.5 | 87.4 - 91.9 | |
| | Stage III | 86.5 - 89.1 | 69.3 - 73.0 | 59.3 - 63.3 | 90.5 - 92.9 | 77.6 - 81.5 | 71.0 - 75.9 | |
| | Stage IV | 45.1 - 49.8 | 16.9 - 20.4 | 9.7 - 12.6 | 49.5 - 54.3 | 19.5 - 23.7 | 11.8 - 15.6 | |
| | Stage UNK | 61.4 - 67.8 | 49.2 - 55.9 | 41.4 - 48.1 | 76.5 - 81.5 | 68.8 - 75.6 | 64.5 - 72.3 | |
| Rectum | Stage I | 95.5 - 98.0 | 88.5 - 92.9 | 80.9 - 87.1 | 97.1 - 100.1 | 94.9 - 100.0 | 92.0 - 99.8 | |
| | Stage II | 90.0 - 94.4 | 76.6 - 83.0 | 67.0 - 74.2 | 92.8 - 97.0 | 82.8 - 89.9 | 76.0 - 85.2 | |
| | Stage III | 91.9 - 94.8 | 75.7 - 80.7 | 62.3 - 68.3 | 92.7 - 96.1 | 78.8 - 84.8 | 67.9 - 75.8 | |
| | Stage IV | 52.2 - 60.7 | 20.7 - 27.8 | 11.6 - 17.6 | 53.2 - 61.5 | 21.6 - 29.1 | 12.2 - 19.0 | |
| | Stage UNK | 80.5 - 87.2 | 64.1 - 72.2 | 55.5 - 64.0 | 85.6 - 91.3 | 74.0 - 82.0 | 67.9 - 77.4 | |
| Lung | Stage I | 85.9 - 88.6 | 64.5 - 68.5 | 50.9 - 55.4 | 90.0 - 92.7 | 74.3 - 78.4 | 63.6 - 68.9 | |
| | Stage II | 67.5 - 73.2 | 41.7 - 47.9 | 31.6 - 37.7 | 74.7 - 80.4 | 50.3 - 58.0 | 41.1 - 49.3 | |
| | Stage III | 48.7 - 52.5 | 20.2 - 23.4 | 13.7 - 16.6 | 54.6 - 59.4 | 26.2 - 31.3 | 20.1 - 25.1 | |
| | Stage IV | 22.1 - 24.1 | 6.7 - 8.0 | 3.9 - 5.0 | 26.2 - 29.3 | 8.4 - 10.7 | 4.9 - 6.7 | |
| | Stage UNK | 31.7 - 39.3 | 17.6 - 23.9 | 13.6 - 19.2 | 42.1 - 53.9 | 30.4 - 43.5 | 25.1 - 37.7 | |
| Prostate | Stage I | 96.4 - 98.0 | 91.4 - 93.9 | 88.0 - 91.0 | 98.1 - 100.2 | 96.5 - 100.5 | 97.1 - 103.2 | |
| | Stage II | 97.3 - 98.1 | 92.1 - 93.4 | 86.9 - 88.6 | 99.9 - 100.7 | 99.8 - 101.4 | 99.7 - 102.4 | |
| | Stage III | 98.5 - 99.5 | 94.4 - 96.5 | 89.0 - 92.0 | 99.6 - 101.7 | 97.5 - 102.1 | 97.2 - 104.3 | |
| | Stage IV | 78.5 - 82.3 | 49.7 - 54.6 | 35.3 - 40.4 | 84.7 - 89.1 | 59.7 - 66.7 | 45.4 - 53.8 | |
| | Stage UNK | 84.2 - 88.6 | 70.1 - 75.3 | 61.6 - 66.6 | 93.4 - 96.3 | 87.9 - 92.7 | 84.9 - 90.4 | |

NOTE: -- Not applicable. If the standard error of a survival rate is between 0.05 and 0.1, the 95% confidence interval of the rate is shown in the table. If it is > 0.1 or is not estimable due to a small number of cases or events then the survival rate is suppressed and denotes as \S . Survival rates include age at diagnosis 15 - 99.

Please refer to the <u>Introduction and Technical Notes</u> for further information.

Data Source: BC Cancer Registry Date Retrieved: 17DEC2019 Analysis by: Cancer Surveillance and Outcomes, Data & Analytics